

LINKS NOT CONCORDANT TO BOUNDARY LINKS

CHARLES LIVINGSTON

(Communicated by Frederick R. Cohen)

ABSTRACT. Casson-Gordon invariants are used to prove that certain links in S^3 are not concordant to boundary links. These examples were first described by Cochran and Orr.

Tim Cochran and Kent Orr recently announced the construction of 2 component links in S^3 with the property that, although all the Milnor $\bar{\mu}$ -invariants vanish, the links are not concordant to boundary links [CO]. They also provide examples of higher dimensional links of two components which are not concordant to boundary links. Their 3-dimensional examples include the links L_m ($m > 0$) illustrated in Figure 1 (see p. 1131). Both bands on the Seifert surface for K_1 are untwisted; that is, the Seifert form vanishes on the homology classes represented by x and y . They let K be the trefoil. The knot K_1 (with $m = 1$) was used in my paper with Gilmer [GL1] as an example of a slice knot that is algebraically doubly null concordant, but not doubly null concordant. The results of [CO] and [GL1] are related. We will show here that the result that L_m is not concordant to a boundary link follows readily from the results of Gilmer [G] on which the work in [GL1] was based. Gilmer's work was based on earlier work on knot concordance by Levine [L] and Casson and Gordon [CG1, CG2].

Danny Ruberman [R1] generalized the work of [GL1] to higher dimensions. (See also [R2] and [Sm] for corrections to [R1].) His observation that the Casson-Gordon invariants apply to higher dimensional slice problems is relevant here. However, a more careful analysis of the Casson-Gordon invariants is needed to prove that the high dimensional examples of [CO] are not concordant to boundary links. That analysis will be presented in a separate paper being written with Pat Gilmer [GL2].

Tim Cochran has informed me that he is also investigating the relationship between the work announced in [CO] and the Casson-Gordon invariant.

The rest of this note is devoted to the proof that L_m is not concordant to a boundary link.

Received by the editors August 30, 1989.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 57M25.

Key words and phrases. Link concordance, boundary links, Casson-Gordon invariant.

©1990 American Mathematical Society
0002-9939/90 \$1.00 + \$.25 per page

Pat Gilmer proved that if a slice knot K in S^3 bounds a genus 1 Seifert surface F , then there is a simple closed curve J on F representing a primitive element in $H_1(F)$ and for which Tristram p -signatures, $\sigma_{(s/p)}(J)$, vanish for all $0 < s < p$, where p is any prime dividing the order of the first homology of the 2-fold cyclic cover of S^3 branched over K . Gilmer produced J as follows. Suppose that K bounds a disk D in B^4 . The surface $F \cup D$ bounds a 3-manifold R in B^4 . Let i denote the inclusion of F into R , and let H denote $i_*^{-1}(\text{Tor}(H_1(R)))$. Gilmer showed that H is a 1-dimensional summand of $H_1(F)$ and that if J represents a generator of H then it has the desired signature properties.

To proceed, consider the link $L = K_1 \cup K_2$. K_1 is clearly slice. (Perform ambient surgery on the unknotted curve y on the evident Seifert surface F_1 .) We will show that if L is concordant to a boundary link then K_1 bounds a disk D in B^4 with the property that $F_1 \cup D$ bounds a 3-manifold R for which the generator of the summand H described above is represented by x .

The construction of D and R is done in two steps. First, suppose that L is concordant in $S^3 \times [0, 1]$ to the boundary link $L' = K'_1 \cup K'_2$ bounding disjoint surfaces F'_1 and F'_2 . Denote the two components of the concordance C_1 and C_2 . The following transversality argument shows that the closed surface $F_1 \cup F'_1 \cup C_1$ bounds a 3-manifold R_1 in $S^3 \times [0, 1]$ with R_1 disjoint from C_2 . (This was stated without proof in [S].)

Let ν denote tubular neighborhood. There is a map p of $(S^3 \times \{0\} - \nu(K_1 \cup K_2)) \cup (S^3 \times \{1\} - \nu(K'_1 \cup K'_2)) \cup \partial(\nu(C_2))$ to S^1 such that p is transverse to 1, $p^{-1}(1) = F_1 \cup F'_1$ and $p^{-1}(-1)$ contains $\partial(\nu(C_2))$. Constructing R_1 via transversality depends on extending p to $S^3 \times [0, 1] - \nu(C_1 \cup C_2)$.

Homotopy classes of maps of a space X to S^1 correspond to elements of $H^1(X)$. The restriction map of

$$H^1(S^3 \times [0, 1] - \nu(C_1 \cup C_2))$$

to

$$H^1((S^3 \times \{0\} - \nu(K_1 \cup K_2)) \cup (S^3 \times \{1\} - \nu(K'_1 \cup K'_2)) \cup \partial(\nu(C_2)))$$

is a map of $\mathbb{Z}^2$ to $\mathbb{Z}^3$ with image consisting of those elements that agree on the meridians of K_1 and K'_1 . Our map p represents a class in the image, and hence p is the restriction of a map as desired.

Let D_1 be a slice disk for K'_1 in B^4 , and let R'_1 be a 3-manifold bounded by $F'_1 \cup D_1$ in B^4 . Form the union of $S^3 \times [0, 1]$ with B^4 identifying $S^3 \times \{1\}$ with ∂B^4 . The union of C_1 with D_1 forms a slice disk D for K_1 . The union of R_1 and R'_1 forms a 3-manifold R bounded by $F_1 \cup D$. Note that R is in the complement of the connected surface $E = C_2 \cup F'_2$ bounded by K_2 .

If an element z in $H_1(F_1)$ represents torsion in $H_1(R)$, it is also torsion in $H_1(B^4 - E)$. The element z can be written as $z = ax + by$. In $H_1(B^4 - E)$,




FIGURE 1

x represents zero and y an element of infinite order, in fact the generator. (Linking numbers of x and y with E in B^4 equal their linking numbers with K_2 in S^3 .) It follows that b is 0, and hence that H is generated by the class represented by x .

The curve x is of the same knot type as K . The order of the homology of the 2-fold cyclic branched cover of S^3 branched over K_1 is $(2m+1)^2$. Hence, as long as $\sigma_{(s/p)}(K) \neq 0$ for some prime divisor p of $(2m+1)$ and $0 < s < p$, L will not be concordant to a boundary link. The trefoil works for all $m > 0$.

REFERENCES

- [CO] T. Cochran and K. Orr, *Not all links are concordant to boundary links*, preprint.
- [CG1] A. Casson and C. McA. Gordon, *Cobordism of classical knots*, preprint, Orsay; reprinted in *A la recherche de la topologie Perdue* (A. Marin and L. Guillou, eds.), Progress in Math. vol. 62, Birkhauser, 1986.
- [CG2] —, *On slice knots in dimension three*, Proc. Sympos. Pure Math. **XXX** (1978), Part two, 39–53.
- [G] P. Gilmer, *Slice knots in S^3* , Quart. J. Math. Oxford **39** (1983), 305–322.
- [GL1] P. Gilmer and C. Livingston, *On embedding 3-manifolds in 4-space*, Topology **22** (1983), 241–252.
- [GL2] —, *The Casson-Gordon invariant and link concordance*, preprint.
- [L] J. Levine, *Knot cobordism groups in codimension two*, Comment. Math. Helv. **44** (1969), 229–244.
- [R1] D. Ruberman, *Doubly slice knots and the Casson-Gordon invariants*, Trans. Amer. Math. Soc. **279** (1983), 569–588.
- [R2] —, *Casson-Gordon invariants and high-dimensional knot theory*, Trans. Amer. Math. Soc. **306** (1988), 579–596.
- [S] N. Sato, *Cobordisms of semi-boundary links*, Topology Appl. **18** (1984), 225–234.
- [Sm] L. Smolinsky, *Casson-Gordon invariants of some 3-fold branched covers of knots*, Topology Appl. **31** (1989), 243–252.