

MAPS OF SURFACE GROUPS TO FINITE GROUPS WITH NO SIMPLE LOOPS IN THE KERNEL

CHARLES LIVINGSTON

*Department of Mathematics
 Indiana University
 Bloomington, IN 47405
 livingst@indiana.edu*

Received 2 April 2000

ABSTRACT

Let F_g denote the closed orientable surface of genus g . What is the least order finite group, G_g , for which there is a homomorphism $\psi: \pi_1(F_g) \rightarrow G_g$ so that no nontrivial simple closed curve on F_g represents an element in $\text{Ker}(\psi)$? For the torus it is easily seen that $G_1 = Z_2 \times Z_2$ suffices. We prove here that G_2 is a group of order 32 and that an upper bound for the order of G_g is given by g^{2g+1} . The previously known upper bound was greater than $2g^{2^{2g}}$.

For any compact surface F there exists a finite group G and a homomorphism $\psi: \pi_1(F) \rightarrow G$ such that no nontrivial element in the kernel of ψ can be represented by a simple closed curve. Such a homomorphism is said to have *nongeometric kernel*. Casson, Gabai, and Skora [1] have each constructed examples of this (see Section 2 for details). The presence of such examples raises a variety of questions relating to the characterization of the finite groups that can occur in this way. This paper addresses the problem of determining the relationship between the genus of F and the order of G . In the case that F is a torus a complete analysis is straightforward. For instance, the natural projection $\psi: \pi_1(F) \rightarrow H_1(F; Z_2) \cong Z_2 \times Z_2$ has nongeometric kernel.

Our first result concerns the genus 2 closed orientable surface, F_2 . Casson's construction yields a group of order 2^{38} . Skora reduced this order considerably by producing a group of order 2^9 . In Section 3 a group of order $2^5 = 32$, G_2 , is constructed for which there is a homomorphism $\psi_2: \pi_1(F_2) \rightarrow G_2$ having nongeometric kernel. In Section 4 it is proved that no such example can be constructed using a group of order less than 32.

The example in Section 3 is generalized to construct examples for arbitrary genus surfaces in Section 5. The order of the groups constructed is quite small compared to previously constructed examples. As the examples directly generalize the minimal genus 2 example, there is the possibility that they are minimal as well.

1. Notation and Conventions

Throughout this paper all surfaces will be closed and orientable. References to basepoints for the fundamental group of a space are omitted. Since the property of being in the kernel of a homomorphism depends only on the conjugacy class of an element, such omissions will not affect the arguments.

By a *simple loop* on a surface we mean an embedding of the circle S^1 .

We will say that a homomorphism $\psi: \pi_1(F) \rightarrow G$ has *geometric kernel* if some nontrivial element in the kernel can be represented by a simple loop. Otherwise ψ has *nongeometric kernel*.

2. Basic Examples

In this section a procedure of Casson is used to construct for each surface F a finite group G and a surjective homomorphism $\psi: \pi_1(F) \rightarrow G$ such that ψ has nongeometric kernel. The orders of the groups involved is computed for contrast with the examples produced in Section 5.

The statement that ψ has nongeometric kernel can be reinterpreted in terms of covering spaces as follows. Corresponding to $\text{Ker}(\psi)$ there is a connected regular covering space $p: \tilde{F} \rightarrow F$ with $p_*(\pi_1(\tilde{F})) = \text{Ker}(\psi)$. An element in $\pi_1(F)$ is in $\text{Ker}(\psi)$ if and only if when represented by a closed path, the path lifts to a closed path in $\tilde{F}$. Hence a simple loop on F represents an element in $\text{Ker}(\psi)$ if and only if it can be lifted to a simple loop in $\tilde{F}$. Conversely, if $p: \tilde{F} \rightarrow F$ is a regular covering space with the property that no nontrivial simple loop on F lifts to $\tilde{F}$ then the natural projection $\psi: \pi_1(F) \rightarrow \pi_1(F)/p_*(\pi_1(\tilde{F}))$ has nongeometric kernel.

Construction Given a surface F , construct the covering space $p: \tilde{F} \rightarrow F$ corresponding to the kernel of the projection $\pi_1(F) \rightarrow H_1(F; \mathbb{Z}_2)$. Simple nonseparating loops on F represent generators of $H_1(F; \mathbb{Z}_2)$ and hence do not lift to $\tilde{F}$. Nontrivial separating simple loops do lift, but each preimage on $\tilde{F}$ is nonseparating on $\tilde{F}$.

Now construct the covering $q: \bar{F} \rightarrow \tilde{F}$ corresponding to the kernel of the projection $\pi_1(\tilde{F}) \rightarrow H_1(\tilde{F}; \mathbb{Z}_2)$. As no nonseparating simple loop on $\tilde{F}$ lifts to $\bar{F}$ it is apparent that no nontrivial simple loop on F lifts to $\bar{F}$.

It remains to show that the covering $p \circ q: \bar{F} \rightarrow F$ is regular; that is, that $p_* \circ q_*(\pi_1(\bar{F}))$ is normal in $\pi_1(F)$. Observe that $q_*(\pi_1(\bar{F}))$ is a characteristic subgroup of $\pi_1(\tilde{F})$ and $p_*(\pi_1(\tilde{F}))$ is a characteristic subgroup of $\pi_1(F)$. Since a characteristic subgroup of a characteristic subgroup is characteristic, $p_* \circ q_*(\pi_1(\bar{F}))$ is characteristic in $\pi_1(F)$, and is hence normal.

Order of $\pi_1(F)/\langle p_* \circ q_*(\pi_1(\bar{F})) \rangle$ The order of this finite group is equal to the degree of the covering $p \circ q$. Suppose that F is of genus g . The Euler characteristic of F is $2 - 2g$. Since $\tilde{F}$ is a 2^{2g} fold cover of F , the Euler characteristic of $\tilde{F}$ is $2^{2g}(2 - 2g)$. The genus of $\tilde{F}$ is $\frac{1}{2}(2 - 2^{2g}(2 - 2g)) = \frac{1}{2}((g - 1)2^{2g+1} + 2) = g'$. The covering $q: \bar{F} \rightarrow \tilde{F}$ is of degree $2^{2g'}$. The degree of $p \circ q: \bar{F} \rightarrow F$ is the product of these two degrees: $2^{2g'}2g = 2^{(g-1)2^{2g+1}+2+2g}$.

Note The construction of Gabai differs considerably from the one above. He notes that every simple curve is in the complement of some index three subgroup of $\pi_1(F)$; nonseparating curves are not in the kernel of some map to Z_3 and separating curves are mapped to a 3-cycle in the third symmetric group, S_3 , under some homomorphism and hence map to the complement of an index three subgroup of S_3 . Hence the quotient map $\pi_1(F) \rightarrow \pi_1(F)/H$ has nongeometric kernel, where H is the intersection of all index three subgroups of $\pi_1(F)$. We have been unable to find reasonable bounds on the size of this quotient.

3. A Small Genus 2 Example

The groups constructed in the previous section are of very large order. If F is of genus 2, the corresponding group is of order 2^{38} . This section presents a description of a group of order 32, G_2 , and a homomorphism ψ_2 of the fundamental group of the genus 2 surface to G_2 , such that ψ_2 has a nongeometric kernel. The next section contains a proof that this example is minimal.

For the remainder of this section F will denote a genus 2 surface.

Construction of G_2 For our purposes, the easiest way to describe G_2 is as follows. Define a group structure on the set $(Z_2)^4 \times Z_2$ by defining the product by

$$(a_1, b_1, a_2, b_2, \epsilon)(a'_1, b'_1, a'_2, b'_2, \epsilon') = (a_1 + a'_1, b_1 + b'_1, a_2 + a'_2, b_2 + b'_2, \epsilon + \epsilon' + b_1 a'_1 + b_2 a'_2).$$

The operations within the parenthesis are addition and multiplication in Z_2 . The verification that this defines a group structure can be done by a direct calculation, which is left to the reader. The group is denoted 32_{42} in [2], and $\Gamma_5 a_1$ in the notation of [3].

An essential calculation for later purposes is that of commutators in G_2 . A direct computation yields

$$[(a_1, b_1, a_2, b_2, \epsilon), (a'_1, b'_1, a'_2, b'_2, \epsilon')] = (0, 0, 0, 0, (b_1 a'_1 - b'_1 a_1) + (b_2 a'_2 - b'_2 a_2)). \quad (1)$$

From this it is apparent that both the center and commutator subgroup of G_2 consists of the set $(0, 0, 0, 0) \times Z_2$. The abelianization of G_2 is $(Z_2)^4$, given by the projection $(Z_2)^4 \times Z_2 \rightarrow (Z_2)^4 \times \{0\}$.

Construction of ψ_2 Let $\{x_1, y_1, x_2, y_2\}$ be a standard generating set of $\pi_1(F)$ so that $\pi_1(F)$ has presentation $\langle x_1, y_1, x_2, y_2, [x_1, y_1][x_2, y_2] = 1 \rangle$. This set projects to a standard symplectic basis of $H_1(F; Z_2)$, $\{|x_1|, |y_1|, |x_2|, |y_2|\}$.

(In the present setting, working with Z_2 coefficients, we do not need to concern ourselves with signs, but in Section 5 sign issues become essential. Note that via a choice of orientation for the surface we can arrange that the algebraic intersection numbers $x_1 \cap y_1$ and $x_2 \cap y_2$ can both be assumed to be -1 . This yields the intersection forms on $H_1(F; Z_2)$ as we use below and on $H_1(F; Z_k)$ as used in Section 5.)

Define $\psi_2: \pi_1(F) \rightarrow G_2$ by setting:

$$\begin{aligned}\psi_2(x_1) &= (1, 0, 0, 0) \times (0), \\ \psi_2(y_1) &= (0, 1, 0, 0) \times (0), \\ \psi_2(x_2) &= (0, 0, 1, 0) \times (0), \\ \psi_2(y_2) &= (0, 0, 0, 1) \times (0).\end{aligned}$$

Using (1) it is easily verified that this gives a well defined surjective representation.

The key observation is that ψ_2 has the following property: if ω_1 and ω_2 are elements of $\pi_1(F)$, then

$$[\psi_2(\omega_1), \psi_2(\omega_2)] = (0, 0, 0, 0) \times (|\omega_1| \cap |\omega_2|), \tag{2}$$

where $|\omega_1| \cap |\omega_2|$ is the Z_2 intersection number of the classes in $H_1(F; Z_2)$ represented by ω_1 and ω_2 . This follows from (1) along with the fact that if (a_1, b_1, a_2, b_2) and (a'_1, b'_1, a'_2, b'_2) are classes in $H_1(F; Z_2)$, then $(a_1, b_1, a_2, b_2) \cap (a'_1, b'_1, a'_2, b'_2) = (b_1 a'_1 - b'_1 a_1) + (b_2 a'_2 - b'_2 a_2)$. (Note also that the natural map $\pi_1(F) \rightarrow H_1(F; Z_2)$ factors through G_2 via ψ_2 .)

The kernel of ψ_2 is nongeometric Suppose that there is a simple loop γ representing a nontrivial element ω in $\text{Ker}(\psi_2)$. Our first observation is that γ can be chosen to be separating. If γ is nonseparating, pick a simple loop γ' meeting γ transversely in exactly one point. Let ω' be the element of $\pi_1(F)$ represented by γ' . Clearly, $[\omega, \omega']$ is in the kernel of ψ_2 and it is represented by a separating simple loop.

Since γ is now assumed to be separating, it bounds a punctured torus on F . This follows from the classification of surfaces. Hence $\omega = [\omega_1, \omega_2]$, where ω_1 and ω_2 are represented by simple loops meeting transversely in one point. From this one computes using (2) that $\psi_2(w) = [\psi_2(\omega_1), \psi_2(\omega_2)] = (0, 0, 0, 0) \times (|\omega_1| \cap |\omega_2|) = (0, 0, 0, 0) \times (1)$, which is nontrivial in G_2 . This contradicts the assumption that $w \in \text{Ker}(\psi_2)$.

4. Minimality of G

The goal of this section is to prove that if F is of genus 2 and the order of G is less than 32, than any homomorphism $\phi: \pi_1(F) \rightarrow G$ has geometric kernel.

Here is a summary of the argument. We first prove that any $\phi: \pi_1(F) \rightarrow G$ has geometric kernel if G is a cyclic extension of an abelian group, that is, if G contains a normal abelian subgroup with cyclic quotient. The approach used to prove this was pointed out by Allan Edmonds. The argument depends on an analysis of the action of the homeomorphism group of F on the set of representations of $\pi_1(F)$ to G . We next note that with the exception of two groups of order 24, $SL_2(Z_3)$ and S_4 , all groups of order less than 32 are cyclic extensions of abelian groups. This can be proved by a case-by-case analysis using the Sylow theorems. More easily, group

tables such as [2] provide the necessary information. The proof is completed using specialized arguments for $SL_2(Z_3)$ and S_4 .

Results of Nielsen and Edmonds We will need results of Nielsen and Edmonds in the arguments that follow. Since, in the original sources, these are stated in terms of group actions as opposed to representations, we will briefly summarize the results now. The papers [4, 5] serve as complete references for this discussion.

In [4, Theorem 1.1] Edmonds proves that two free actions of a finite abelian group G on a closed orientable surface are equivalent via an equivariant, orientation-preserving homeomorphism if and only if they determine the same class in $\Omega_2^{free}(G)$, the bordism group of free G actions on surfaces. It follows from the bordism spectral sequence (see [6] for a more complete discussion than in [4, 5]) that $\Omega_2^{free}(G) = H_2(G, Z)$. For G cyclic, $H_2(G, Z) = 0$, so as a corollary one has the result of Nielsen [7] that any two orientation-preserving free actions of a cyclic group on a closed orientable surface are equivalent via an equivariant homeomorphism.

In [5] the case of finite nonabelian split metacyclic groups is addressed. Here the main theorem of the introduction implies that two such actions are equivalent if they determine the same element in $H_2(G, Z)$. In the case of G a dihedral group, D_{2p} , of order $2p$, p odd, it is shown (see Lemma 1.2 of [5]) that $H_2(G, Z) = 0$. Hence, as in the cyclic case, any two such actions are equivalent. We will use this in the case that G is the third symmetric group, $S_3 = D_6$.

We will need the translation of these results to the setting of representations of surface groups to finite groups. Given a free action of G on M , there is a quotient surface N along with a surjective representation $\pi_1(N) \rightarrow G$. It follows from covering space theory (see Lemmas 2.1 and 2.2 of [4]) that there is a bijection between equivalence classes of free actions of G on M and equivalence classes of surjective representations $\pi_1(N) \rightarrow G$. Initially this equivalence of representations is defined via the action of the group of automorphisms of $\pi_1(N)$, but via a theorem of Nielsen all such automorphisms are induced by (basepoint preserving) homeomorphisms of the surface. That is, two representations ϕ and ϕ' are equivalent if there is a homeomorphism h of N such that $\phi' = \phi \circ h_*$. Notice that ϕ has geometric kernel if and only if $\phi \circ h_*$ has geometric kernel.

Summarizing, we have the following.

Lemma 1. *If G is either a finite cyclic group or $G = S_3$, then any two surjective representations of $\pi_1(F)$ to G are equivalent via the action of the homeomorphism group.*

Cyclic Extension of Abelian Groups

Since, with the two exceptions dealt with below, all groups of order less than 32 contain normal abelian subgroups with cyclic quotient, we begin by observing the following consequence of Lemma 1.

Lemma 2. *If G is a cyclic group and $\phi: \pi(F) \rightarrow G$ is a surjective homomorphism, then there is a homeomorphism h of F such that $\phi \circ h_*(x_1)$ generates G ,*

and $\phi \circ h_*(y_1)$, $\phi \circ h_*(x_2)$ and $\phi \circ h_*(y_2)$ are all trivial.

Proof. Let ϕ' be a representation for which $\phi'(x_1)$ generates G and $\phi'(y_1)$, $\phi'(x_2)$ and $\phi'(y_2)$ are all trivial in G . Then according to Lemma 1, ϕ and ϕ' are equivalent, so there is a homeomorphism h as desired $\square$.

Theorem 3. *If G contains an abelian normal subgroup N such that G/N is cyclic, then any surjective homomorphism $\phi: \pi_1(F) \rightarrow G$ has a geometric kernel.*

Proof. Denote the quotient map $G \rightarrow G/N$ by ρ . Applying Lemma 2 we can assume that $\rho \circ \phi(x_2)$ and $\rho \circ \phi(y_2)$ are both trivial. Hence $\phi(x_2)$ and $\phi(y_2)$ are both in N . The commutator $[x_2, y_2]$ is represented by a simple loop and is in the kernel of ϕ , since $\phi([x_2, y_2])$ is in the commutator subgroup of an abelian group.

Exceptional Groups

Case 1 We begin by recalling that $SL_2(Z_3)$ is isomorphic to the semidirect product of the quaternionic 8-group, Q , with Z_3 . We will use the standard notation for elements in Q . The generator of Z_3 will be denoted t . The action of Z_3 on Q is given by $tit^{-1} = j$, $tjt^{-1} = k$ and $tk t^{-1} = i$. Note that $t(-1)t^{-1} = -1$ and that $-1 \in Q$ is hence central in $SL_2(Z_3)$.

Suppose $\phi: \pi_1(F) \rightarrow SL_2(Z_3)$ is a surjective representation with nongeometric kernel. Applying Lemma 2 to the composition $\pi_1(F) \rightarrow SL_2(Z_3) \rightarrow SL_2(Z_3)/Q = Z_3$ we can assume that $\phi(x_1) = tq_1$, $\phi(y_1) = q_2$, $\phi(x_2) = q_3$, and $\phi(y_2) = q_4$, where each q_i is in Q .

Since $[x_2, y_2]$ is represented by a simple loop, $[q_3, q_4] \neq 1$. Hence $[q_3, q_4] = -1 \in Q$. It follows that $[tq_1, q_2] = -1$. Note that $q_2 \neq \pm 1$, so $q_2 = \pm i, \pm j$, or $\pm k$. From the commutator relation, $tq_1q_2q_1^{-1}t^{-1}q_2^{-1} = -1$, we have that $tq_1q_2q_1^{-1}t^{-1} = -q_2$. Given that $q_2 \neq \pm 1$ it follows that $-q_2 = q_2^{-1}$ and so $tq_1q_2q_1^{-1}t^{-1} = q_2^{-1}$. For any two quaternions, $q_1q_2q_1^{-1} = q_2^{\pm 1}$. Hence, $tq_2t^{-1} = q_2^{\pm 1}$. However, this is impossible, given that $q_2 \neq \pm 1$ and the action of t on Q .

Case 2 The symmetric group S_4 is the semidirect product of $Z_2 \times Z_2$ with S_3 . As a subgroup, the $Z_2 \times Z_2$ is given by the set $\{(1), (12)(34), (13)(24), (14)(23)\}$. The subset $\{(1), (12), (13), (23), (123), (321)\}$ yields the $S_3 \subset S_4$.

Let $\phi: \pi_1(F) \rightarrow S_4$ be a surjective representation with nongeometric kernel. Lemma 1 applied to the composition $\pi_1(F) \rightarrow S_4 \rightarrow S_4/(Z_2 \times Z_2) \cong S_3$ shows that by applying a homeomorphism we can arrange that ϕ takes on the values $\phi(x_1) = (12)n_1$, $\phi(y_1) = n_2$, $\phi(x_2) = (123)n_3$, and $\phi(y_2) = n_4$, where each n_i is in $Z_2 \times Z_2$. (The representation $\rho: \pi_1(F) \rightarrow S_3$ defined by $\rho(x_1) = (12)$, $\rho(y_1) = 1$, $\rho(x_2) = (123)$, and $\rho(y_2) = 1$ must, according to Lemma 1, be equivalent to the composition $\pi_1(F) \rightarrow S_4 \rightarrow S_4/(Z_2 \times Z_2) \cong S_3$, and hence we can apply a homeomorphism to achieve the desired form.)

Since both y_1 and y_2 are represented by simple loops, neither n_2 nor n_4 are trivial. Also, $[x_1, y_1]$ is represented by a simple loop, so $[(12)n_1, n_2] \neq 1$. It follows that $n_2 \neq (12)(34)$, since $(12)(34)$ commutes with n_1 and with (12) . There are two

other possibilities for n_2 .

Suppose that $n_2 = (13)(24)$. There are three possible values of n_4 to be considered. Because y_1y_2 is realized by a simple loop, $n_4 \neq (13)(24)$. It is easily seen that $y_2x_1y_1^{-1}x_1^{-1}$ is realized by a simple loop. Hence $n_4 \neq (12)(34)$. Finally $n_4 \neq (14)(23)$, because $y_1x_2y_2^{-1}x_2^{-1}$ can also be represented by a simple loop.

We proceed similarly if $n_2 = (14)(23)$. Clearly $n_4 \neq (14)(23)$. Because $y_1x_2^{-1}y_2^{-1}x_2$ is realized by a simple loop, $n_4 \neq (13)(24)$. Finally, it is again easily seen that $y_2x_1y_1^{-1}x_1^{-1}$ is realized by a simple loop. This implies that $n_4 \neq (12)(34)$. All possibilities have now been eliminated $\square$.

5. Generalizations

The group constructed in Section 3, G_2 , can be generalized to a group G_k such that for the genus k surface F_k there is a homomorphism $\phi_k : \pi_1(F_k) \rightarrow G_k$ with nongeometric kernel. The arguments are similar to those of Section 3 and are only outlined here.

Define G_k by defining a product on the set $(Z_k)^{2k} \times Z_k$ as follows.

$$(a_1, b_1, a_2, b_2, \dots, b_k, \epsilon)(a'_1, b'_1, a'_2, b'_2, \dots, b'_k, \epsilon') = \\ (a_1 + a'_1, b_1 + b'_1, a_2 + a'_2, b_2 + b'_2, \dots, b_k + b'_k, \epsilon + \epsilon' + \sum b_i a'_i)$$

Sums and products within the parenthesis are in Z_k . That this defines a group is a straightforward calculation.

There is a natural representation $\phi_k : \pi_1(F_k) \rightarrow G_k$ as before. In this case the essential observation is

$$[\phi_k(\omega_1), \phi_k(\omega_2)] = (0, 0, \dots, 0) \times (|\omega_1| \cap |\omega_2|) \quad (1)$$

where $(|\omega_1| \cap |\omega_2|)$ is the Z_k intersection number of the classes in $H_1(F_k; Z_k)$ represented by ω_1 and ω_2 .

If ϕ_k had geometric kernel, there would be a separating simple loop representing an element in the kernel. Using the classification of surfaces, that element would be of the form

$$[\omega_1, \omega'_1][\omega_2, \omega'_2] \cdots [\omega_m, \omega'_m]$$

with $m < k$ and $(|\omega_i| \cap |\omega'_i|) = 1$ for all i . A contradiction follows as in Section 3 $\square$.

Remark The order of the group G just constructed is g^{2g+1} . This number should be contrasted to the order found in Section 2, $2^{(g-1)2^{2g+1}+2+2g}$. The first is obviously much smaller than the second. The results of this paper, along with our difficulties in trying to find smaller examples, leads us to conjecture that g^{2g+1} represents the least possible order.

6. Addenda

Recently Pikaart [8] has announced a proof that for g odd the least order of a group G for which there is a map $\pi_1(F) \rightarrow G$ with nongeometric kernel is g^{2g+1} (as we conjecture) if one adds the constraint that the kernel be a characteristic subgroup, not just normal. That this may be a nontrivial constraint follows from Pikaart's theorem for g even, in which case he shows that the smallest *characteristic* quotient is of order $2^{2g}g^{2g+1}$, which is much larger than our normal quotients. Pikaart's results depend on the results of this paper.

Acknowledgements

Thanks are due to Allan Edmonds for pointing out the proof of Theorem 3. The work in Section 5 was motivated by discussions with Dennis Johnson. Special thanks are due to the referee for carefully reading this paper and contributing to its clarity and correctness.

References

- [1] R. Skora, *Dissertation*, Department of Mathematics, University of Texas, Austin, Texas, 1984.
- [2] A. D. Thomas and G. V. Wood, *Group Tables*, Shiva Publishing Limited, Kent, Great Britain, 1980.
- [3] M. Hall and J. K. Senior, *The groups of order 2^n ($n \leq 6$)* Macmillan, New York (1964).
- [4] A. Edmonds, *Surface Symmetry I*, Michigan J. Math **29** (1982) 171–183.
- [5] A. Edmonds, *Surface Symmetry II*, Michigan J. Math **30** (1983) 143–154.
- [6] P. Conner and E. Floyd, *Differentiable Periodic Maps*, Springer, New York, 1964.
- [7] J. Nielson, *Die Struktur periodischer Transformationen von Flächen*, Dansk Vid. Selsk., Mat.-Fys. Medd. **15** (1937), 1–77.
- [8] M. Pikaart, *Large characteristic subgroups of surface groups not containing any simple loops*, preprint posted at the Mathematics ArXiv: <http://front.math.ucdavis.edu/math.GT/0005093>.